

DR ELTON'S FOOT POST OPERATIVE INSTRUCTIONS

Incision care

- Keep the post-operative dressings **on** until your first follow-up appointment in 5-7 days. (you may shower with the dressings on using garbage bags)
- At your first follow-up appointment we will change your bandages and reapply steri-strips. Your sutures will be removed at your second appointment 10-14 days after surgery.
- There is no need to apply Neosporin or other ointments to the incision.
- Please alert us to any pus drainage or redness around the incision that is painful to the touch.

Activity

- You may not place weight on your leg for 24 hours after surgery. Once the numbing medicine has worn off, you may weight bear on your heel with a protective shoe. In some cases, you will be heel weight bearing for 6 weeks after surgery. You will need crutches or a walker to go home with after surgery. If you need crutch training we can have you see the physical therapist before surgery.
- If you have been given a fracture boot, please bring this to your first follow-up appointment.
- Elevate your operative leg whenever possible. Elevation means placing your leg above the level of your heart.
- Icing with ice packs or using an ice compression device will significantly help reduce your swelling and pain.
- If you need physical therapy, it will likely not begin until 2-6 weeks after surgery. We will decide at your first post-operative when physical therapy begins.
- Please wear the white compressive stocking (TED hose) on your non-operative leg until your first post-operative visit. These are helpful in preventing blood clots and reducing swelling. They may be removed for showering and laundering.
- Please ask about driving at your first visit, we will make recommendations on an individual basis. We recommend no driving while you are taking narcotic pain medicine or while your driving leg is in a splint or boot.

Medications

- We recommend you pick up your prescriptions prior to surgery. Do not take them before surgery unless directed.
- You will receive prescriptions for pain medicines; take these only if you need them. Usually your medication will not contain Tylenol so you may supplement with Tylenol three times a day unless we advise you otherwise.
- In most cases, you will be prescribed a blood thinner for 2 weeks. Start taking it the morning <u>after</u> surgery.
- While taking narcotic pain medicine, two common reactions are possible.
 - Constipation is the most common side effect. Maintain a diet high in fiber and use any over the counter stool softeners (Colace, Metamucil, Fiber-One, etc.). Drinking plenty of water will help too.
 - Secondly, some people may experience itching or mild rashes with narcotic pain medicine. This can be treated with over-the-counter diphenhydramine (Benadryl) per the package instructions. Breathing problems associated with pain medicine usually require immediate evaluation in the emergency room.
- **Plan appropriately for refill request**. Refills will not be authorized after office hours or on the weekend or holidays. Per our office policy, we have 24 hours to authorize a refill request.
- Please do not take over the counter anti-inflammatories (ibuprofen or Aleve) unless we have told you otherwise.

Follow up appointments

If a follow up appointment was not scheduled for you prior to surgery, please call the office to schedule. We would like to see you 5-7 days after surgery unless we have told you otherwise. We typically see you again at 10-14 days and remove your sutures. Additional appointments will be at 6 weeks, 12 weeks, and 6 months after surgery.

Emergencies

• Call our office immediately with any questions or concerns. If you feel the emergency is life threatening, go to the nearest emergency room. Due to Dr. Elton and Kelli's clinic and surgery schedules, we will make every attempt to call you back before the end of the day; however, if your need is urgent someone on our staff will assist you until we can be in direct contact with you. For urgent after hour matters, there is a PA on call that can assist you.